



## **Sucrose acetate isobutyrate based nanogels as liquid fiducial tissue markers with potential use in image guided radiotherapy**

**Bruun, Linda Maria; Schaarup-Jensen, Henrik; Jølck, Rasmus Irming; Hansen, Anders Elias; Christiansen, Anders N.; Clausen, Mads Hartvig; Kjær, Andreas; Scherman, Per Jonas Bengtsson; Andresen, Thomas Lars**

*Publication date:*  
2014

[Link back to DTU Orbit](#)

### *Citation (APA):*

Bruun, L. M., Schaarup-Jensen, H., Jølck, R. I., Hansen, A. E., Christiansen, A. N., Clausen, M. H., Kjær, A., Scherman, P. J. B., & Andresen, T. L. (2014). *Sucrose acetate isobutyrate based nanogels as liquid fiducial tissue markers with potential use in image guided radiotherapy*. Abstract from Liposome Research Days 2014, Copenhagen, Denmark.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## SUCROSE ACETATE ISOBUTYRATE BASED NANOGELS AS LIQUID FIDUCIAL TISSUE MARKERS WITH POTENTIAL USE IN IMAGE GUIDED RADIOTHERAPY

Linda M. Bruun<sup>1,2</sup>, Henrik Schaarup-Jensen<sup>1,3</sup>, Rasmus I. Jølck<sup>1,2</sup>, Anders E. Hansen<sup>1,2</sup>, Anders N. Christiansen<sup>4</sup>, Mads H. Clausen<sup>1,3</sup>, Andreas Kjær<sup>5</sup>, Per Jonas Bengtsson Scherman<sup>6</sup> and Thomas L. Andresen<sup>1,2</sup>

Technical University of Denmark, Center for Nanomedicine and Theranostics<sup>1</sup>, 2800 Kgs. Lyngby, Denmark

Technical University of Denmark, Department of Micro-and Nanotechnology<sup>2</sup>, 2800 Kgs. Lyngby, Denmark

Technical University of Denmark, Department of Chemistry<sup>3</sup>, 2800 Kgs. Lyngby, Denmark

Technical University of Denmark, DTU Compute, Department of Applied Mathematics and Computer Science<sup>4</sup>, 2800 Kgs. Lyngby, Denmark

Rigshospitalet and University of Copenhagen, Department of Clinical Physiology, Nuclear Medicine & PET and Cluster for Molecular Imaging<sup>5</sup>, 2100 Copenhagen, Denmark.

Department of Oncology, Section of Radiotherapy<sup>6</sup>, Rigshospitalet, 2100 Copenhagen, Denmark.

[linbr@nanotech.dtu.dk](mailto:linbr@nanotech.dtu.dk), [hensaar@kemi.dtu.dk](mailto:hensaar@kemi.dtu.dk)

### SUMMARY

Sucrose Acetate IsoButyrate (SAIB)-based radiopaque fiducial tissue markers containing coated gold nanoparticles (AuNPs) have been developed. One of these formulations, the PNIPAM (poly(N-isopropyl acrylamide))-AuNP-SAIB based gel, was assessed to be a suitable marker for image guided radiotherapy (IGRT).

### INTRODUCTION

IGRT is an important tool used to visualize tumors during radiation treatment. Tumors rarely display a fixed position during irradiation or within a treatment period due to breathing motion, changes in organ filling and tumor size. Radiopaque fiducial tissue markers are therefore of great interest to place within - or around tumors in order to achieve high precision in IGRT. Better markers will enable improved tumor coverage i.e. the destruction of all cancer tissue. Furthermore, it is important to have a safer and more accurate setup for tumor treatment in radiotherapy - including adaptation of online "tumor-tracking" - thereby delivering high radiation doses to tumors with minimal damage to surrounding healthy tissue [1].


### EXPERIMENTAL METHODS

The poster presents the development of novel injectable liquid fiducial markers based on formulations of SAIB and polylactide (PLA) containing coated AuNPs. A mixture of SAIB/EtOH/PLA (75:20:5) was used as matrix due to its high biocompatibility, low viscosity, as well as its ability to form a stable, biodegradable gel depot upon injection [2]. Three different AuNP coating systems were tested - a dithiolane functionalized SAIB derivative, synthesized in four steps starting from sucrose, together with PEG- and PNIPAM polymers, respectively. The AuNPs were synthesized by a three step seeding protocol using chloroauric acid as Au<sup>3+</sup> source and trisodium citrate as reductant and stabilizer. *In vitro* release from SAIB gels was studied in PBS-buffer at 37°C. Gels (200µL) containing SAIB/EtOH/PLA (75:20:5) + 30mg PNIPAM-AuNPs mL<sup>-1</sup> or 10mg PEG-AuNPs mL<sup>-1</sup> were

injected subcutaneously at the upper left flank of immunocompetent NMRI-mice using hypodermic 25G needles. The gel-depots were visualized by micro-CT imaging.

### RESULTS AND DISCUSSION

The dithiolane functionalized SAIB derivative was discarded as a coating option due to irreversible aggregation of the formed nanoparticles. The SAIB gels containing PEGylated AuNPs provided high CT contrast *in vivo*, however it suffered from AuNP migration towards the gel-boundaries resulting in an inhomogeneous distribution of contrast agent. The PNIPAM-coated-AuNP-SAIB gel provided both excellent CT contrast and high *in vivo* stability. Furthermore, the PNIPAM-coated-AuNPs are easy to handle, as they can be lyophilized and stored as an air stable powder, which is readily dispersible in ethanol.



Maximum Intensity Projection of PNIPAM-AuNP-SAIB gel depot in a mouse viewed from several angles.

### CONCLUSION

The PNIPAM-AuNP-SAIB gel showed high stability, biocompatibility and provided high CT contrast *in vivo*. This gel is a viable alternative to existing liquid fiducial markers.

### REFERENCES

1. Engstrom P.E., Haraldsson P., Landberg T., et al., *Acta Oncol.* **44**(6), 572-578 (2005)
2. Jølck, R. I., Binderup T., Hansen A.E., et al. *Adv. Healthcare Mater.* (2014).



**Liposome Research Days**  
Living Innovation 4-7 August 2014

